

The further development of out-of-office BP monitoring: Japan's ImPACT Program Project's achievements, impact, and direction

1 | INTRODUCTION

New guidelines for the management of hypertension were recently released by the European Society of Hypertension/European Society of Cardiology (the 2018 ESH/ESC guidelines) and the American College of Cardiology/American Heart Association (the 2017 ACC/AHA guidelines).^{1,2} These guidelines stress the importance of out-of-office blood pressure (BP) values over that of office BP values. In Japan and other Asian countries, we have

highlighted the use of the out-of-office BP-guided management of hypertension.³⁻⁵ Ambulatory BP monitoring (ABPM) and home BP monitoring (HBPM) are the two standard measurements of out-of-office BP. Their use can detect masked (uncontrolled) hypertension (normotension in office BP and hypertension in out-of-office BP) in individuals at the highest risk of cardiovascular events.⁶⁻⁸ There are three clinical phenotypes of masked hypertension: morning hypertension, daytime hypertension, and nocturnal hypertension.⁹




FIGURE 1 Abnormal 24-h ambulatory BP profiles and actisensitivity in a 72-y-old woman who developed B-type aortic dissection during the daytime. A, The 24-h trend of ambulatory BP and the parameters measured by IMS-ABPM. Upper: The riser pattern was detected in summer (8 mo before onset). Lower: An excessive morning surge was detected in autumn (4 mo before onset). B, Actisensitivity calculated as the slope of daytime ambulatory BP change against the physical activity change. The red plots and the regression line represent the data of 72-y-old woman who developed aortic dissection. Upper: The inverse actisensitivity was found in summer. Lower: The actisensitivity was within the normal range




FIGURE 2 A novel approach to individual management for hypertension. Biologic and environmental signals corresponding to an individual's living conditions at home were simultaneously collected from multiple devices and sensors by an IoT (Internet of Things) gateway and bridges based on the hybrid Wi-SUN/Wi-Fi transmission system. This high-speed analysis and reporting system facilitates hypertensive management. (http://www.jichi.ac.jp/usr/card/research/index_en.html). BLE, Bluetooth®

Ambulatory BP monitoring has traditionally been considered the gold standard to detect the risk of high BP throughout the 24-hour day, and its use can detect all three types of masked hypertension. Toward the achievement of the goal of “zero” cardiovascular events, three components are needed for “perfect 24-hour BP control”: (a) lowering the 24-hour BP level, (b) maintaining an adequate circadian rhythm, and (c) avoiding excessive BP variability including the morning BP surge.¹⁰ All three of these components can be assessed by ABPM. Extremely disrupted patterns of circadian rhythm of nighttime BP and exaggerated morning BP surge such as the riser pattern (higher nighttime BP than daytime BP) and the extreme-dipper pattern (excessive nighttime BP falls) are reported to be associated with cardiovascular risk.^{11–13}

Home BP monitoring is frequently used in clinical practice and to identify masked hypertension defined by self-measured home BP. The use of HBPM can detect the risk of morning hypertension.^{14–18} The recently developed “nighttime HBPM” automatically obtains and records BP values at fixed intervals while an individual is sleeping, and it can be used as an alternative to ABPM for the assessment of nighttime BP.¹³ Two additional modalities have been developed to detect the risk of hypertension during sleep (especially in patients with obstructive sleep apnea): (a) “trigger nighttime HBPM” with a

hypoxic episode-trigger function and a heart rate-trigger function and (b) beat-by-beat continuous surge BP monitoring.^{13,19}

Home BP monitoring can thus detect the risk of morning hypertension and that of nocturnal hypertension. However, HBPM would underestimate the risk of daytime hypertension, because HBPM measures an individual's blood pressure in the less stressful resting condition at home. A patient's behavior, surrounding environment, and various triggering factors affect his or her daytime ambulatory BP changes. Masked daytime hypertension (ie, normotension in office BP and hypertension in daytime BP) induced by physical activity or work- or home-related psychological stress can be detected only by ABPM.

2 | NEW MULTI-SENSOR ABPM AND BP SENSITIVITY INDEXES

We recently developed a device that provides information/communication technology (ICT)-based multi-sensor ABPM (IMS-ABPM), which can store all of the waveforms of intra-cuff pressures during oscillometric BP measurement.²⁰ One of the limitations of the original ABPM is the accuracy of “real” daytime BP measurements, because the daytime movement of the ABPM device wearer's upper arm may




FIGURE 3 Time-trend reporting system of ambulatory BP and environmental conditions along with an individual's high-risk places and times

modify the intra-cuff pressure and produce BP reading artifacts. By excluding the abnormal BP values with abnormal waveforms as artifacts, we can evaluate the “real” daytime BP measurements more accurately. In addition, based on the IMS-ABPM's stored waveforms and its function of detecting an irregular heartbeat (IHB),²¹ the device can be used for the screening of atrial fibrillation (AF).

The IMS-ABPM device is equipped with a thermosensor, a highly sensitive actigraph, and an atmospheric pressure sensor to simultaneously assess the triggers of BP surge (a pressor component of BP variability).²⁰ By determining the association between many subjects' BP measurements and these triggers, new BP sensitivity indexes could be calculated; for example, the slope of the ambulatory BP values against specific triggers such as temperature (thermosensitivity evaluation), physical activity (actisensitivity), atmospheric pressure (atmospheric sensitivity), humidity (humidity sensitivity), and more. Based on these new indicators, we could classify the characteristics of hypertension with excessive BP sensitivity to specific triggers such as thermosensitive hypertension, actisensitive hypertension, atmospheric hypertension, and humidity-sensitive hypertension. These sensitivities might overlap and augment each other.

In fact, the actisensitivity of BP (the slope of daytime BP change against physical activity change during a 5-minute period before the BP measurement) is augmented in the cold winter season compared to the warm summer in the same patients.²² This may partly explain the winter increase in the rate of cardiovascular events. The

IMS-ABPM device can specifically detect masked daytime hypertension with a physical activity-induced BP surge.

As an example: a 72-year-old woman developed B-type aortic dissection during the daytime. Although she had been treated with amlodipine, candesartan, and hydrochlorothiazide, the IMS-ABPM device detected abnormal 24-hour BP profiles during the year prior to the onset of aortic dissection. The riser pattern was detected in the summer (8 months before the onset), and an excessive morning surge was detected in the autumn (4 months before the onset) (Figure 1A). The patient's simultaneously calculated actisensitivity was also disrupted. The actisensitivity was within the normal range in autumn; however, the inverse actisensitivity (physical activity reduced the ambulatory BP) was found in summer (Figure 1B). The disrupted BP regulation against physical activity, which is modified by environmental (eg, seasonal) conditions, might produce the excessive BP surge that triggers cardiovascular events.

3 | THE IMPACT HYBRID WI-SUN/WIFI TRANSMISSION SYSTEM

Based on recent technology developments, it has been suggested that the use of ICT-based devices and a real-time feedback IoT (Internet of Things)-based system could facilitate a novel approach to patient management.^{22,23} In the ImPACT program (IMpulsing PARadigm Change




FIGURE 4 Time series out-of-office BP-based anticipation management of hypertension: Effective for bridging the gap between the guidelines and individualized medicine

through disruptive Technologies program of the Cabinet Office, Government of Japan), we have successfully developed an integrated system that collects both biologic and environmental data, with the hybrid Wi-SUN/Wi-Fi transmission system (Figure 2). Environmental sensors introduced in a patient's home continuously monitor the temperature, humidity, and illumination in different rooms with different conditions throughout 24-hour periods. An ICT-based wrist-type pulse wave monitoring device (wearable monitoring) which we also developed in the ImPACT program continuously monitors the wearer's pulse rate, pulse wave, and activity.

With the combination of these biological sensor devices, environmental sensors, and the hybrid Wi-SUN/Wi-Fi transmission system, the environmental determinants of BP surges (ie, the pressor component of BP variability) with different time phases could be used to identify the riskiest places and times at which an individual's maximum BP surge is exaggerated at home throughout the year (Figure 3). Health information technology (HIT) solutions like this are increasingly being recognized as an important component of the advances in health care, and the latest version of the ACC/AHA hypertension guidelines highlight the importance and emerging roles of HIT.¹

4 | BLOOD PRESSURE SURGE-BASED ANTICIPATION MEDICINE

Using these new technologies, it is hoped that the occurrence of cardiovascular events could be anticipated based on data obtained by these novel approaches to out-of-office patient monitoring, with the ultimate goal of eliminating the occurrence of cardiovascular events in patients with hypertension. Such an approach is referred to as "anticipation medicine" for zero cardiovascular events, within which BP variability is a key biomarker.^{10,19,20,22}

The self-monitoring of blood pressure using HBPM, particularly when combined with telemonitoring, has recently been shown to facilitate the titration of antihypertensive therapy in subjects with poorly controlled hypertension in general practice, without increasing the general practitioner's workload.^{24,25} This highlights the potential for ICT-based out-of-office BP measurement solutions in clinical practice.

There is a gap between the guideline-initiated general management of hypertension and the individualized optimal management of hypertension. An increase in the number of out-of-office BP measurements could increase the sensitivity and specificity of the average BP-based diagnosis of hypertension (as guideline-based medicine), and this increase in data could also detect the various specific trigger-induced BP surges, that is, diurnal, day-by-day, and seasonal BP surges (as individualized medicine) (Figure 4). The resonance hypothesis holds that the exaggerated pathological surge BP generated by the resonance of different BP surges with different time phases would trigger cardiovascular events.²⁶ The times and places at which the surge BP is generated would be the most risky times/places for an individual's cardiovascular event onset. Even among well-controlled hypertensive patients, the morning BP surge remains significant in the winter.²⁷ Anticipation medicine for cardiovascular diseases—which both anticipates pathological surge BP based on the previous time series of individual BP data and avoids the generation of surge BP—is an ideal future practical direction to take in order to decrease the gap between the guidelines and individualized medicine in the era of ICT-based "real-world" big data analysis and feedback systems.

ACKNOWLEDGMENTS

This paper was supported in part by the IMPulsing PARadigm Change through disruptive Technologies (ImPACT) program of the Cabinet

Office, Government of Japan. We thank Makoto Kato, Hiroko Masaki, Tomohide Sato, Satoshi Hoshide, Tomoyuki Kabutoya, Kimiyo Saito, and Tomoko Shiga for their support. We also thank Shinobu Ozaki, Yoshiteru Nozoe, Shinichi Takahashi, and Takahiro Fujiwara from A&D Co. for developing IMS-ABPM, wearable wrist pulse monitoring, and the ICT-based data collecting system, and Takeya Shigezumi, Terumi Sata, and Takashi Naiki from Rohm Co. for developing multi-sensors and the real-time and hybrid Wi-SUN/Wi-Fi transmission system.

CONFLICT OF INTEREST

This paper was supported in part by the Impulsing PARadigm Change through disruptive Technologies (IMPACT) program of the Cabinet Office, Government of Japan. Nobuhiko Yasui is an employee of A&D Co. Other authors report no conflicts of interest to disclose.

ORCID

Kazuomi Kario  <https://orcid.org/0000-0002-8251-4480>

Kazuomi Kario MD, PhD¹ 

Naoko Tomitani BSc¹

Hiroshi Kanegae BSc²

Nobuhiko Yasui BSE³

Ryozo Nagai MD, PhD⁴

Hiroshi Harada PhD⁵

¹Department of Medicine, Division of Cardiovascular Medicine, Jichi Medical University School of Medicine (JMU), Shimotsuke, Japan

Email: kkario@jichi.ac.jp

²Genki Plaza Medical Center for Health Care, Tokyo, Japan

³A&D Co, Tokyo, Japan

⁴Jichi Medical University School of Medicine, Shimotsuke, Japan

⁵Graduate School of Informatics, Kyoto University, Kyoto, Japan

Correspondence

Kazuomi Kario, Department of Medicine, Division of Cardiovascular Medicine, Jichi Medical University School of Medicine, Shimotsuke, Japan.

Email: kkario@jichi.ac.jp

REFERENCES

- Whelton PK, Carey RM, Aronow WS, et al. 2017 ACC/AHA/AAPA/ABC/ACPM/AGS/APhA/ASH/ASPC/NMA/PCNA guideline for the prevention, detection, evaluation, and management of high blood pressure in adults: executive summary: a report of the American College of Cardiology/American Heart Association Task Force on Clinical Practice Guidelines. *J Am Coll Cardiol*. 2018;71:2199-2269.
- Williams B, Mancia G, Spiering W, et al. 2018 ESC/ESH guidelines for the management of arterial hypertension. *Eur Heart J*. 2018;39:3021-3104. [Epub ahead of print]
- Shimamoto K, Ando K, Fujita T, et al. The Japanese Society of Hypertension guidelines for the management of hypertension (JSH 2014). *Hypertens Res*. 2014;37:253-390.
- Kario K, Park S, Buranakitjaroen P, et al. Guidance on home blood pressure monitoring: a statement of the HOPE Asia Network. *J Clin Hypertens*. 2018;20:456-461.
- Kario K, Tomitani N, Buranakitjaroen P, et al. Home blood pressure control status in 2017-2018 for hypertension specialist centers in Asia: Results of the AsiaBP@Home study. *J Clin Hypertens*. 2018;20:1686-1695. In press
- Fujiwara T, Yano Y, Hoshide S, Kanegae H, Kario K. Association of cardiovascular outcomes with masked hypertension defined by home blood pressure monitoring in a Japanese general practice population. *JAMA Cardiol*. 2018;3:583-590.
- Pierdomenico SD, Pierdomenico AM, Coccina F, et al. Prognostic value of masked uncontrolled hypertension. *Hypertension*. 2018;72(4):862-869.
- Kario K, Chen CH, Park S, et al. Consensus document on improving hypertension management in Asian patients, taking into account Asian characteristics. *Hypertension*. 2018;71:375-382.
- Kario K. Global Impact of 2017 American Heart Association/American College of Cardiology Hypertension Guidelines: a perspective from Japan. *Circulation*. 2018;137:543-545.
- Kario K. *Essential Manual on Perfect 24-hour Blood Pressure Management from Morning to Nocturnal Hypertension: Up-to-date for Anticipation Medicine*. Tokyo, Japan: Wiley-Blackwell; 2018:1-309.
- Kario K, Pickering TG, Umeda Y, et al. Morning surge in blood pressure as a predictor of silent and clinical cerebrovascular disease in elderly hypertensives: a prospective study. *Circulation*. 2003;107:1401-1406.
- Kario K, Pickering TG, Matsuo T, Hoshide S, Schwartz JE, Shimada K. Stroke prognosis and abnormal nocturnal blood pressure falls in older hypertensives. *Hypertension*. 2001;38:852-857.
- Kario K. Nocturnal hypertension: new technology and evidence. *Hypertension*. 2018;71:997-1009.
- Hoshide S, Yano Y, Haimoto H, et al. Morning and evening home blood pressure and risks of incident stroke and coronary artery disease in the Japanese general practice population: The Japan Morning Surge-Home Blood Pressure Study. *Hypertension*. 2016;68:54-61.
- Ohkubo T, Imai Y, Tsuji I, et al. Home blood pressure measurement has a stronger predictive power for mortality than does screening blood pressure measurement: a population-based observation in Ohasama, Japan. *J Hypertens*. 1998;16:971-975.
- Kario K, Saito I, Kushiro T, et al. Home blood pressure and cardiovascular outcomes in patients during antihypertensive therapy: Primary results of HONEST, a large-scale prospective, real-world observational study. *Hypertension*. 2014;64:989-996.
- Kario K, Saito I, Kushiro T, et al. morning home blood pressure is a strong predictor of coronary artery disease: The HONEST Study. *J Am Coll Cardiol*. 2016;67:1519-1527.
- Kario K, Iwashita M, Okuda Y, et al. Morning home blood pressure and cardiovascular events in Japanese hypertensive patients. *Hypertension*. 2018;72:854-861.
- Kario K. Evidence and Perspectives on the 24-hour management of hypertension: hemodynamic biomarker-initiated 'anticipation medicine' for zero cardiovascular event. *Prog Cardiovasc Dis*. 2016;59:262-281.
- Kario K. Hemodynamic biomarker-initiated anticipation medicine in the future management of hypertension. *Am J Hypertens*. 2017;30:226-228.
- Kabutoya T, Imai Y, Hoshide S, Kario K. Diagnostic accuracy of a new algorithm to detect atrial fibrillation in a home blood pressure monitor. *J Clin Hypertens (Greenwich)*. 2017;19:1143-1147.
- Kario K, Tomitani N, Kanegae H, et al. Development of a new ICT-based multisensor blood pressure monitoring system for use in hemodynamic biomarker-initiated anticipation medicine for

- cardiovascular disease: The National IMPACT Program Project. *Prog Cardiovasc Dis*. 2017;60:435-449.
23. Impulsing Paradigm Change through Disruptive Technologies Program (ImPACT): "An Ultra Big Data Platform for Reducing Social Risks." Predictive health care realized by using ultra-big data with connected medical devices [movie: MP4]. http://www.jichi.ac.jp/usr/card/research/index_en.html. Accessed December 26, 2018.
24. Nishizawa M, Hoshide S, Okawara Y, Matsuo T, Kario K. Strict blood pressure control achieved using an ICT-based home blood pressure monitoring system in a catastrophically damaged area after a disaster. *J Clin Hypertens (Greenwich)*. 2017;19(1):26-29.
25. McManus RJ, Mant J, Franssen M, et al. Efficacy of self-monitored blood pressure, with or without telemonitoring, for titration of antihypertensive medication (TASMINH4): an unmasked randomised controlled trial. *Lancet*. 2018;391:949-959.
26. Kario K. New insight of morning blood pressure surge into the triggers of cardiovascular disease – synergistic resonance of blood pressure variability. *Am J Hypertens*. 2016;29:14-16.
27. Nishizawa M, Fujiwara T, Hoshide S, et al. Winter morning surge in blood pressure after the Great East Japan Earthquake. *J Clin Hypertens*. 2019;21:208-216